



TABLE I: Oligodeoxynucleotides

A. Thermodynamics and Kinetics

14-dC-S	ATG CAG CTA AGT CA
14-dC-A	TGA CTT AGC TGC AT
14-BrdC-S	ATG CAG CTA AGT CA
14-BrdC-A	TGA CTT AGC TGC AT
14-MedC-S	ATG <u>CAG</u> <u>CTA</u> AGT <u>CA</u>
14-MedC-A	TGA <u>CTT</u> <u>AGC</u> <u>TGC</u> <u>AT</u>
12-dC-S	CAT GCA GCC CCA
12-dC-A	TGG GGC TGC ATG
12-BrdC-S	CAT GCA GCC CCA
12-MedC-S	<u>CAT</u> <u>GCA</u> <u>GCC</u> <u>CCA</u>
16-dC-A	TGG GGC TGC ATG GCG T
16-BrdC-S	ACG CCA TGC AGC CCC A

Nomenclature: Oligodeoxynucleotide length - (dC, MedC, BrdC) - (S = sense; A = antisense).
C = 5'-Bromodeoxycytidine, C = 5'-Methyldeoxycytidine.

B. Linkers, Displacers and Targets

PstI

GAT GAT GAT GTG CAG CCA ATG CCC CAG GAG CCC T	P-D-MedC
GAT GAT GAT GTG CAG CCA AAG CCC CAG GAG CCC T	P-D-BrdC-E(10)
GAT GAT GAT GTG CAG CCA ATG CCC CAG GAG CCC A	P-D-BrdC-E(24)
GAT GAT GAT GTG CAG CCA ATG CCC CAG GAG CCC T	P-D-BrdC
G TAC CTA CTA CTA C	P-L-dC
*G CCA ATG CCC CAG GAG CCC T	P-T-D
AC GTC GGT TAC GGG GTC CTC GGG A	P-T-L

EcoRI

CCT CGA AGG AGC CTT CCA CAG CCG AAT TGT AGT AGT AGT AAG CT	E-D-BrdC
CA TCA TCA TCA T	E-L-dC
*CCT CGA AGG AGC CTT CCA CAG CCG	E-T-D
GGA GCT TCC TCG GAA GGT GTC GGC TTA A	E-T-L

XmaI (SmaI)

TCT CGG CTC ACT GCA ACG TCC GCC TCC CGG GTA GTA GTA GTA	X-D-BrdC
CAT CAT CAT CAT	X-L-dC
CC CAT CAT CAT CAT	Sm-L-dC
*TCT CGG CTC ACT GCA ACG TCC GCC TC	X-T-D
AGA GCC GAG TGA CGT TGC AGG CGG AGG GCC	X-T-L

SfiI

GTA GTA GTA GTA CCC CGG CCA CAC ACA CAC ACA CAC GA	S-D-BrdC
CAT CAT CAT CAT	S-L-dC

NotI

CTC TCT CTC TCT CTC TCT GCG GCC GTA GTA GTA GTA	N-D-BrdC
CAT CAT CAT CAT	N-L-dC

Nomenclature: Oligodeoxynucleotides are named x-y-z where:
x = restriction site (first letter only)
y = function (displacer, linker or target)
z = composition (BrdC, MedC, dC) or polarity (displacer or linker side)
E(N) = error at position N (a mismatch) if present

PMS-ES
PMS-SE

PMS-NH
PMS-HN

Nomenclature: Oligodeoxynucleotides are named x-y-z where x = BT if both branch migration and triplex formation are possible, x = B0 if only branch migration is possible; y = composition (dC, MedC or BrdC) in the branch migration region (1) and/or triplex forming region (2); z = laboratory index.



TABLE II: Bromodeoxycytidine and Methyldeoxycytidine Thermodynamics

	t_m in 1 M Na ⁺ , C = 6 μ M		
Oligodeoxynucleotides	pH 4	pH 7	pH 10
14-dC-A + 14-dC-S		57°	
14-dC-A + 14-BrdC-S		63°	
14-dC-S + 14-BrdC-A		62.5°	
14-BrdC-A + 14-BrdC-S		65°	
12-dC-A + 12-dC-S	60°	53.5°	50°
12-dC-A + 12-BrdC-S	70°	69.5°	54°
	pH 4.7	pH 7	pH 9.6
14-dC-A + 14-dC-S		57°	
14-dC-A + 14-MedC-S		60°	
14-dC-S + 14-MedC-A		59.5°	
14-MedC-A + 14-MedC-S		64.5°	
12-dC-A + 12-dC-S	50°	60°	47.5°
12-dC-A + 12-MedC-S	55°	67°	50.5°

[illegible]



TABLE III: Temperature Dependence of Displacement Rates with BrdC-Containing Displacers

A. Blunt ends

Oligodeoxynucleotides	Temp (°C)	Half-time (min) for displacement with 12-BrdC-S at:		
		101 μ M	20 μ M	4 μ M
12-dC-S* (C = 0.25 μ M) +	37	2	4-8	16-32
12-dC-A (C = 0.75 μ M)	32	4-8	8-16	32
	27	4-8	32-64	128-256

B. Overhangs

Oligodeoxynucleotides	(°C)	Half-time (min) for displacement with 16-BrdC-S at:	
		3 μ M	0.57 μ M
12-dC-S* (C = 0.25 μ M) +	37	<1	4-8
16-dC-A (C = 0.57 μ M)	32	<1	16-32
	27	<1	2-4
* 5'- ³² P-labeled	22	<1	<1

C. Effect of Linkers

Site	Overhang	G+C%	Half time for displacement (minutes)	
			No linker	Linker
<i>Eco</i> RI	5'	0	60	6
<i>Pst</i> I	3'	50	8	< 1
<i>Xma</i> I	5'	100	8-16	< 1

CONDITIONS: Ligase buffer (pH = 7); 37°C; 10 μ L per reaction.
Target: Kinased strand 10 ng, Unlabeled strand 30 ng.
Displacer: 150 ng; Linker (if present): large molar excess.
Displacer concentration = 1 μ M.

0926660



* by definition

Displacer	Relative yield (Fragment B/Fragment A)	
	Experimental	Calculated
P-D-BrdC	24	29
P-D-BrdC-E(10)	1	1
P-D-BrdC-E(24)	24	39

United States Patent & Trademark Office

Office of Initial Patent Examination


Application papers not suitable for publication

SN 09 898 750

Mail Date 07-03-01

- ☐ Non-English Specification
- ☐ Specification contains drawing(s) on page(s) _____ or table(s) _____
- ☐ Landscape orientation of text ☐ Specification ☐ Claims ☐ Abstract
- ☐ Handwritten ☐ Specification ☐ Claims ☐ Abstract
- ☐ More than one column ☐ Specification ☐ Claims ☐ Abstract
- ☐ Improper line spacing ☐ Specification ☐ Claims ☐ Abstract
- ☒ Claims not on separate page(s)
- ☐ Abstract not on separate page(s)
- ☐ Improper paper size -- Must be either A4 (21 cm x 29.7 cm) or 8-1/2"x 11"
- ☐ Specification page(s) _____ ☐ Abstract
- ☐ Drawing page(s) _____ ☐ Claim(s)
- ☐ Improper margins
- ☐ Specification page(s) _____ ☐ Abstract
- ☐ Drawing page(s) _____ ☐ Claim(s)
- ☐ Not reproducible
- | <u>Reason</u> | <u>Section</u> |
|---|--|
| <input type="checkbox"/> Paper too thin | <input type="checkbox"/> Specification page(s) _____ |
| <input type="checkbox"/> Glossy pages | <input type="checkbox"/> Drawing page(s) _____ |
| <input type="checkbox"/> Non-white background | <input type="checkbox"/> Abstract |
| | <input type="checkbox"/> Claim(s) |
- ☐ Drawing objection(s)
- ☐ Missing lead lines, drawing(s) _____
- ☐ Line quality is too light, drawing(s) _____
- ☐ More than 1 drawing and not numbered correctly
- ☐ Non-English text, drawing(s) _____
- ☐ Excessive text, drawing(s) _____
- ☐ Photographs capable of illustration, drawing(s) _____

United States Patent & Trademark Office
Office of Initial Patent Examination -- Scanning Division



Application deficiencies found during scanning:

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

☐ Page(s) _____ of _____ were not present
for scanning. (Document title)

✓ Scanned copy is best available. *Drawings*